MADISON NEUROLOGY · NEUROLOGY SYMPTOM GUIDE

Acting out dreams or injuring oneself or a bed partner during sleep

Structured history, interpretation & clinical review

Symptom 90 of 94
Clinical educational workbook
Version 1.0 · October 2026

Not for clinical use. This guide cannot diagnose, triage, or exclude an emergency. Seek emergency care for severe, sudden, or worsening symptoms.

Contents

Scope and safe use 3

Section 1: Input form 4

1A. Immediate safety screen 4

1B. Timeline 4

1C. Discriminating features 4

1D. Associated findings 5

1E. Clinical context 5

Section 2: Interpreting the input form 7

Section 3: Clinical review 9

Selected references 11

> **Document purpose**
> A structured educational history and diagnostic-reasoning workbook. Treatment is outside its scope.

Scope and safe use

Acting out dreams or causing sleep-related injury is characterized by dream recall, timing, violent/complex behavior, medication, age, and Parkinsonism, cognitive, or autonomic clues.

Not for clinical use.
This educational workbook does not diagnose, triage, estimate probability, replace examination, or exclude an emergency. If a safety-screen answer is Yes or uncertain, follow the stated action rather than continuing routine interpretation.

How to complete the workbook

- Record the current episode and the person's usual baseline; use a witness account for events with impaired awareness.
- Check one option per question unless a clinician documents a different combination. "Not sure" is unresolved information, not a negative answer.
- Complete the safety screen before all other sections. Record exact onset or last-known-well time for sudden deficits.
- Use Section 2 only after the form is complete. More than one pattern may be relevant; urgency takes precedence.
- Use Section 3 to plan focused examination and clinician-selected investigations. No test is automatic or universally required.

Specific escalation rule

New severe headache, new focal weakness or numbness, new speech/language difficulty, reduced consciousness, a prolonged seizure, severe breathing or swallowing difficulty, sudden visual loss, or rapidly progressive weakness requires emergency assessment. Call 911 in the United States.

Section 1: Input form

Checkbox history for Acting out dreams or injuring oneself or a bed partner during sleep

1A. Immediate safety screen

Is there current severe breathlessness, blue lips, chest pain, or inability to awaken?

☐ No ☐ Yes ☐ Not sure

If Yes: Call emergency services now.

Is the person falling asleep while driving, operating machinery, or doing another safety-critical activity?

☐ No ☐ Yes ☐ Not sure

If Yes: Stop the activity now and arrange prompt assessment; do not drive until the safety issue has been evaluated.

Is there new focal weakness, sudden severe headache, confusion, or a prolonged seizure?

☐ No ☐ Yes ☐ Not sure

If Yes: Call emergency services now.

Has a sleep behavior caused serious injury, or is there a new prolonged unresponsive episode?

☐ No ☐ Yes ☐ Not sure

If Yes: Seek emergency or same-day assessment depending on current responsiveness and injury severity.

> **Stop rule:** Any Yes answer triggers the action printed under that question. Any Not sure answer requires prompt in-person clarification; do not treat it as No.

1B. Timeline

How did the symptom begin?

☐ Suddenly, over seconds or minutes ☐ Over hours to days ☐ Gradually over weeks or months ☐ Longstanding or since childhood ☐ Not sure

Use the first occurrence for recurrent episodes; use the current episode for a changing symptom.

What is the overall course?

☐ One persistent episode ☐ Separate recurrent episodes ☐ Steadily worsening ☐ Fluctuating or variable ☐ Currently resolved

Has this exact pattern been assessed before?

☐ Definitely new ☐ The same pattern occurred before ☐ Present and stable for a long time ☐ Not sure

1C. Discriminating features

Is there a vivid dream matching the behavior?

☐ Yes, often with clear recall on awakening ☐ No dream recall ☐ Not sure

When during sleep does it occur?

☐ Mostly later in the night ☐ Mostly soon after falling asleep ☐ Variable throughout ☐ Not sure

Has the person or bed partner been injured or at risk?

☐ Yes ☐ Near injury or dangerous objects involved ☐ No injury risk ☐ Not sure

Is there Parkinsonism, cognitive change, orthostatic symptoms, constipation, smell loss, or a new antidepressant/other medicine?

☐ Neurological or autonomic clues ☐ Relevant medicine change ☐ Both ☐ None

1D. Associated findings

Are there other neurological symptoms—weakness, numbness, speech change, double vision, swallowing difficulty, or severe imbalance?

☐ Yes ☐ No ☐ Not sure

Are there systemic clues—fever, weight loss, cancer, infection risk, pregnancy/postpartum status, or a new medication?

☐ Yes ☐ No ☐ Not sure

How much does this affect daily activity?

☐ Noticeable but activities continue ☐ Some activities are limited ☐ Cannot perform usual activities ☐ Not sure

1E. Clinical context

Who is being assessed?

☐ Adult or adolescent ☐ Child under 12 ☐ Adult aged 65 or older

Could a medicine, substance, withdrawal, or recent dose change be relevant?

☐ No plausible contributor ☐ Yes or possibly ☐ Not reviewed

Include sedatives, dopamine-blocking drugs, stimulants, decongestants, anticholinergics, alcohol, and non-prescribed substances when relevant.

Is there a relevant neurological, muscular, metabolic, cardiovascular, autoimmune, or sleep disorder?

☐ No known relevant condition ☐ Yes ☐ Not sure

What prior information is available?

☐ A previous examination or investigation ☐ Prior records but no clear explanation ☐ No useful prior assessment ☐ Not sure

Additional history and documentation

Record exact dates/times, witness details, relevant medication doses and changes, prior investigation reports, and the person’s baseline function.

Section 2: Interpreting the input form

Answer combinations generate a differential and an assessment pathway—not a diagnosis.

> **Precedence:** Emergency safety answers override every pattern below. Otherwise, the highest-urgency fully matched pattern takes precedence. If entries are incomplete, contradictory, or match no pattern, use the unresolved-information pathway.

Pattern 1: Typical dream-enactment pattern

Is there a vivid dream matching the behavior? Yes, often with clear recall on awakening
When during sleep does it occur? Mostly later in the night

Prompt clinical review pattern — Promptly

Dream-linked later-night behavior follows a REM-sleep behavior assessment with neurological review.

Differential to review:

- REM-sleep behavior disorder

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Pattern 2: Neurodegenerative-associated dream enactment

Is there Parkinsonism, cognitive change, orthostatic symptoms, constipation, smell loss, or a new antidepressant/other medicine? Neurological or autonomic clues / Both
Is there a vivid dream matching the behavior? Yes, often with clear recall on awakening

Prompt clinical review pattern — Promptly

Dream enactment with movement, cognitive, or autonomic clues needs neurological and sleep assessment.

Differential to review:

- REM-sleep behavior disorder

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Pattern 3: Non-REM or unclear-recall pattern

Is there a vivid dream matching the behavior? No dream recall
When during sleep does it occur? Mostly soon after falling asleep

Prompt clinical review pattern — Promptly

Early behavior without dream recall follows a non-REM parasomnia and seizure-mimic assessment.

Differential to review:

- Non-REM parasomnia or confusional arousal
- Nocturnal seizure or another behavioral mimic

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Pattern 4: Medication-associated enactment

Is there Parkinsonism, cognitive change, orthostatic symptoms, constipation, smell loss, or a new antidepressant/other medicine? Relevant medicine change

Prompt clinical review pattern — Promptly

A relevant drug change requires medication and sleep-breathing review.

Differential to review:

- Medication-associated or sleep-apnea-related enactment

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Unresolved-information pathway: The entries need structured in-person characterization.

Prompt clinical review pattern — Promptly

Do not guess missing answers or convert uncertainty into reassurance. Document what is unknown, perform an in-person examination, clarify the timeline with a witness or records when available, and reassess if any new feature appears.

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Section 3: Clinical review

Diagnostic modules and focused investigation logic

The examination comes before a test list. Select investigations only after localization, urgency, comorbidity, pregnancy status, renal function, anticoagulation, and prior results have been considered.

3.1 REM-sleep behavior disorder

Discriminating features: Dream enactment with vivid recall, later-night timing, complex/violent behavior, and possible neurodegenerative association.

Logical investigations: Sleep-specialist assessment with polysomnography and neurological examination; immediate environmental injury precautions.

3.2 Non-REM parasomnia or confusional arousal

Discriminating features: Early-night behavior without clear dream recall and with incomplete awakening or amnesia.

Logical investigations: Sleep history, trigger review, and video-polysomnography when distinction is uncertain.

3.3 Medication-associated or sleep-apnea-related enactment

Discriminating features: New antidepressant or other medicine, or apnea-related arousals with apparent dream behavior.

Logical investigations: Medication and breathing review with polysomnography as clinically indicated.

3.4 Nocturnal seizure or another behavioral mimic

Discriminating features: Stereotyped frequent episodes, postictal features, or events inconsistent with normal dream behavior.

Logical investigations: Neurological assessment with EEG/video-polysomnography when seizure features are present.

Reassessment and documentation

- If the symptom pattern changes, repeat the safety screen and timeline rather than carrying forward an earlier interpretation.
- If a test is unrevealing but the deficit persists or progresses, revisit localization and seek specialist review; a normal result does not automatically end assessment.
- If symptoms resolve, document duration, completeness of recovery, recurrence, and any residual deficit. Resolved sudden focal symptoms can still require urgent vascular assessment.
- If the history and examination conflict, obtain collateral information and in-person review rather than forcing a single diagnostic label.

Selected references

Sources provide background standards and do not replace local emergency pathways or current specialist guidance. Accessed October 2026.

1. National Institute for Health and Care Excellence. (2021, updated 2024). *Obstructive sleep apnoea/hypopnoea syndrome and obesity hypoventilation syndrome in over 16s (NG202)*. https://www.nice.org.uk/guidance/ng202
2. Winkelman JW, Berkowski JA, DelRosso LM, et al.. (2025). *Treatment of restless legs syndrome and periodic limb movement disorder: an American Academy of Sleep Medicine clinical practice guideline*. https://pubmed.ncbi.nlm.nih.gov/39324694/

Madison Neurology LLC — Neurology Symptom Guide
Acting out dreams or injuring oneself or a bed partner during sleep
Educational reference only. Not for clinical use. Treatment is outside the scope of this workbook.