MADISON NEUROLOGY · NEUROLOGY SYMPTOM GUIDE

Difficulty swallowing

Structured history, interpretation & clinical review

Symptom 53 of 94
Clinical educational workbook
Version 1.0 · October 2026

Not for clinical use. This guide cannot diagnose, triage, or exclude an emergency. Seek emergency care for severe, sudden, or worsening symptoms.

Contents

Scope and safe use 3

Section 1: Input form 4

1A. Immediate safety screen 4

1B. Timeline 4

1C. Discriminating features 4

1D. Associated findings 5

1E. Clinical context 5

Section 2: Interpreting the input form 7

Section 3: Clinical review 9

Selected references 11

> **Document purpose**
> A structured educational history and diagnostic-reasoning workbook. Treatment is outside its scope.

Scope and safe use

Swallowing, chewing, and saliva-control problems are separated into oropharyngeal versus esophageal symptoms, neurological bulbar dysfunction, obstruction, and aspiration risk.

Not for clinical use.
This educational workbook does not diagnose, triage, estimate probability, replace examination, or exclude an emergency. If a safety-screen answer is Yes or uncertain, follow the stated action rather than continuing routine interpretation.

How to complete the workbook

- Record the current episode and the person’s usual baseline; use a witness account for events with impaired awareness.
- Check one option per question unless a clinician documents a different combination. “Not sure” is unresolved information, not a negative answer.
- Complete the safety screen before all other sections. Record exact onset or last-known-well time for sudden deficits.
- Use Section 2 only after the form is complete. More than one pattern may be relevant; urgency takes precedence.
- Use Section 3 to plan focused examination and clinician-selected investigations. No test is automatic or universally required.

Specific escalation rule

New severe headache, new focal weakness or numbness, new speech/language difficulty, reduced consciousness, a prolonged seizure, severe breathing or swallowing difficulty, sudden visual loss, or rapidly progressive weakness requires emergency assessment. Call 911 in the United States.

Section 1: Input form

Checkbox history for Difficulty swallowing

1A. Immediate safety screen

Is there new face droop, one-sided weakness or numbness, difficulty speaking or understanding, double vision, or severe loss of coordination?

☐ No ☐ Yes ☐ Not sure

If Yes: Call emergency services now and record the exact time the person was last known well.

Is there new confusion, reduced consciousness, a seizure, or difficulty awakening?

☐ No ☐ Yes ☐ Not sure

If Yes: Call emergency services now. Do not leave the person alone or give food, drink, or medication by mouth.

Is there new severe breathlessness, inability to swallow saliva, blue lips, or rapidly worsening weakness?

☐ No ☐ Yes ☐ Not sure

If Yes: Call emergency services now; breathing and swallowing problems can become critical quickly.

Is there sudden loss of vision, a new missing visual field, or a painful red eye?

☐ No ☐ Yes ☐ Not sure

If Yes: Seek emergency eye or emergency-department assessment now.

> **Stop rule:** Any Yes answer triggers the action printed under that question. Any Not sure answer requires prompt in-person clarification; do not treat it as No.

1B. Timeline

How did the symptom begin?

☐ Suddenly, over seconds or minutes ☐ Over hours to days ☐ Gradually over weeks or months ☐ Longstanding or since childhood ☐ Not sure

Use the first occurrence for recurrent episodes; use the current episode for a changing symptom.

What is the overall course?

☐ One persistent episode ☐ Separate recurrent episodes ☐ Steadily worsening ☐ Fluctuating or variable ☐ Currently resolved

Has this exact pattern been assessed before?

☐ Definitely new ☐ The same pattern occurred before ☐ Present and stable for a long time ☐ Not sure

1C. Discriminating features

What happens during swallowing?

☐ Difficulty initiating or controlling food/saliva ☐ Coughing, choking, or wet voice during/after swallow ☐ Food sticks in throat or chest ☐ Nasal regurgitation or chewing weakness

Which consistencies are affected?

☐ Liquids mainly ☐ Solids mainly ☐ Both liquids and solids ☐ Saliva control only

Is the problem intermittent, progressive, or suddenly acquired?

☐ Sudden onset ☐ Progressive ☐ Intermittent ☐ Longstanding stable

Is there weight loss, dehydration, recurrent chest infection, or food avoidance?

☐ Yes ☐ No ☐ Not sure

1D. Associated findings

Are there other neurological symptoms—weakness, numbness, speech change, double vision, swallowing difficulty, or severe imbalance?

☐ Yes ☐ No ☐ Not sure

Are there systemic clues—fever, weight loss, cancer, infection risk, pregnancy/postpartum status, or a new medication?

☐ Yes ☐ No ☐ Not sure

How much does this affect daily activity?

☐ Noticeable but activities continue ☐ Some activities are limited ☐ Cannot perform usual activities ☐ Not sure

1E. Clinical context

Who is being assessed?

☐ Adult or adolescent ☐ Child under 12 ☐ Adult aged 65 or older

Could a medicine, substance, withdrawal, or recent dose change be relevant?

☐ No plausible contributor ☐ Yes or possibly ☐ Not reviewed

Include sedatives, dopamine-blocking drugs, stimulants, decongestants, anticholinergics, alcohol, and non-prescribed substances when relevant.

Is there a relevant neurological, muscular, metabolic, cardiovascular, autoimmune, or sleep disorder?

☐ No known relevant condition ☐ Yes ☐ Not sure

What prior information is available?

☐ A previous examination or investigation ☐ Prior records but no clear explanation ☐ No useful prior assessment ☐ Not sure

Additional history and documentation

Record exact dates/times, witness details, relevant medication doses and changes, prior investigation reports, and the person’s baseline function.

Section 2: Interpreting the input form

Answer combinations generate a differential and an assessment pathway—not a diagnosis.

> **Precedence:** Emergency safety answers override every pattern below. Otherwise, the highest-urgency fully matched pattern takes precedence. If entries are incomplete, contradictory, or match no pattern, use the unresolved-information pathway.

Pattern 1: Sudden neurogenic swallowing difficulty

Is the problem intermittent, progressive, or suddenly acquired? Sudden onset
What happens during swallowing? Difficulty initiating or controlling food/saliva / Coughing, choking, or wet voice during/after swallow / Nasal regurgitation or chewing weakness

Emergency warning pattern — Now

Sudden oropharyngeal swallowing difficulty may reflect stroke or acute bulbar dysfunction and requires emergency assessment.

Differential to review:

- Oropharyngeal neurogenic dysphagia or aspiration
- Myasthenia, motor neuron, myopathy, or cranial neuropathy

Action: Call 911 or go to the nearest emergency department now. Do not drive yourself.

Pattern 2: Aspiration or nutritional-risk pattern

Is there weight loss, dehydration, recurrent chest infection, or food avoidance? Yes
What happens during swallowing? Coughing, choking, or wet voice during/after swallow / Difficulty initiating or controlling food/saliva / Nasal regurgitation or chewing weakness

Urgent assessment pattern — Same day

Coughing/aspiration clues with weight loss, dehydration, or chest infections need urgent swallow evaluation.

Differential to review:

- Oropharyngeal neurogenic dysphagia or aspiration

Action: Arrange same-day medical assessment. If symptoms are rapidly worsening or severe, call 911.

Contingency: Until assessed, do not force oral intake; inability to swallow saliva or breathe comfortably is an emergency.

Pattern 3: Mechanical sticking or esophageal pattern

What happens during swallowing? Food sticks in throat or chest
Which consistencies are affected? Solids mainly / Both liquids and solids

Prompt clinical review pattern — Promptly

Food sticking, especially progressive solid-predominant symptoms, follows an esophageal structural/motility pathway.

Differential to review:

- Esophageal obstruction, motility disorder, or reflux-related disease

Action: Arrange an in-person medical or neurological review promptly, usually within days. Seek sooner care if the pattern changes.

Pattern 4: Fatigable or progressive bulbar pattern

What happens during swallowing? Nasal regurgitation or chewing weakness
Is the problem intermittent, progressive, or suddenly acquired? Progressive

Urgent assessment pattern — Same day

Progressive chewing/nasal regurgitation requires prompt neuromuscular bulbar assessment.

Differential to review:

- Myasthenia, motor neuron, myopathy, or cranial neuropathy

Action: Arrange same-day medical assessment. If symptoms are rapidly worsening or severe, call 911.

Unresolved-information pathway: The entries need structured in-person characterization.

Urgent assessment pattern — Same day

Do not guess missing answers or convert uncertainty into reassurance. Document what is unknown, perform an in-person examination, clarify the timeline with a witness or records when available, and reassess if any new feature appears.

Action: Arrange same-day medical assessment. If symptoms are rapidly worsening or severe, call 911.

Section 3: Clinical review

Diagnostic modules and focused investigation logic

The examination comes before a test list. Select investigations only after localization, urgency, comorbidity, pregnancy status, renal function, anticoagulation, and prior results have been considered.

3.1 Oropharyngeal neurogenic dysphagia or aspiration

Discriminating features: Initiation difficulty, coughing/wet voice, nasal regurgitation, weak cough, or stroke/Parkinson/neuromuscular context.

Logical investigations: Urgent swallow and cranial-nerve assessment; speech-language evaluation and instrumental swallow study when indicated.

3.2 Esophageal obstruction, motility disorder, or reflux-related disease

Discriminating features: Food sticking, solid-predominant symptoms, or progressive solid-to-liquid difficulty.

Logical investigations: Gastroenterology assessment and endoscopy/imaging or motility testing as indicated.

3.3 Myasthenia, motor neuron, myopathy, or cranial neuropathy

Discriminating features: Fatigable chewing/swallowing, nasal speech, progressive bulbar signs, or respiratory weakness.

Logical investigations: Bulbar/respiratory examination, junction or electrodiagnostic testing and targeted imaging.

3.4 Structural head/neck, post-surgical, medication, or functional swallowing problem

Discriminating features: Local surgery/radiation, mass, xerostomia, drug effects, or functional variability.

Logical investigations: Oral/head-neck examination, medication review, and instrumental swallow evaluation based on phase.

Reassessment and documentation

- If the symptom pattern changes, repeat the safety screen and timeline rather than carrying forward an earlier interpretation.
- If a test is unrevealing but the deficit persists or progresses, revisit localization and seek specialist review; a normal result does not automatically end assessment.
- If symptoms resolve, document duration, completeness of recovery, recurrence, and any residual deficit. Resolved sudden focal symptoms can still require urgent vascular assessment.
- If the history and examination conflict, obtain collateral information and in-person review rather than forcing a single diagnostic label.

Selected references

Sources provide background standards and do not replace local emergency pathways or current specialist guidance. Accessed October 2026.

1. National Institute for Health and Care Excellence. (2019, updated 2023). *Suspected neurological conditions: recognition and referral (NG127)*. https://www.nice.org.uk/guidance/ng127